

Compactness by compensation and nonlinear parabolic type equation

Marin Mišur^a and Darko Mitrović^b

^aUniversity of Zagreb, Croatia, pfobos@gmail.com

^bUniversity of Montenegro, Montenegro, matematika@t-com.me



Panov's result

The most general version of the classical L^2 results has recently been proved by E. Yu. Panov (2011): Assume that the sequence (\mathbf{u}_n) is bounded in $L^p(\mathbf{R}^d; \mathbf{R}^r)$, $2 \leq p < \infty$, and converges weakly in $\mathcal{D}'(\mathbf{R}^d)$ to a vector function \mathbf{u} .

Let $q = p'$ if $p < \infty$, and $q > 1$ if $p = \infty$. Assume that the sequence

$$\sum_{k=1}^{\nu} \partial_k(\mathbf{A}^k \mathbf{u}_n) + \sum_{k,l=\nu+1}^d \partial_{kl}(\mathbf{B}^{kl} \mathbf{u}_n)$$

is precompact in the anisotropic Sobolev space $W_{loc}^{-1,-2;q}(\mathbf{R}^d; \mathbf{R}^m)$, where $m \times r$ matrices \mathbf{A}^k and \mathbf{B}^{kl} have variable coefficients belonging to $L^{2\bar{q}}(\mathbf{R}^d)$, $\bar{q} = \frac{p}{p-2}$ if $p > 2$, and to the space $C(\mathbf{R}^d)$ if $p = 2$.

We introduce the set $\Lambda(\mathbf{x})$

$$\Lambda(\mathbf{x}) = \left\{ \lambda \in \mathbf{C}^r \mid (\exists \xi \in \mathbf{R}^d \setminus \{0\}) : \left(i \sum_{k=1}^{\nu} \xi_k \mathbf{A}^k(\mathbf{x}) - 2\pi \sum_{k,l=\nu+1}^d \xi_k \xi_l \mathbf{B}^{kl}(\mathbf{x}) \right) \lambda = \mathbf{0}_m \right\},$$

and consider the bilinear form on \mathbf{C}^r

$$q(\mathbf{x}, \lambda, \eta) = \mathbf{Q}(\mathbf{x}) \lambda \cdot \eta, \quad (1)$$

where $\mathbf{Q} \in L_{loc}^{\bar{q}}(\mathbf{R}^d; \text{Sym}_r)$ if $p > 2$ and $\mathbf{Q} \in C(\mathbf{R}^d; \text{Sym}_r)$ if $p = 2$.

Finally, let $q(\mathbf{x}, \mathbf{u}_n, \mathbf{u}_n) \rightharpoonup \omega$ weakly in the space of distributions.

The following theorem holds

Theorem. Assume that $(\forall \lambda \in \Lambda(\mathbf{x})) q(\mathbf{x}, \lambda, \lambda) \geq 0$ (a.e. $\mathbf{x} \in \mathbf{R}^d$) and $\mathbf{u}_n \rightharpoonup \mathbf{u}$, then $q(\mathbf{x}, \mathbf{u}(\mathbf{x}), \mathbf{u}(\mathbf{x})) \leq \omega$.

The connection between q and Λ given in the previous theorem, we shall call *the consistency condition*.

Goal: to formulate and extend the results from the preceding theorem to the $L^p - L^q$ framework for appropriate (greater than one) indices p and q where $p < 2$.

Localisation principle

For $\alpha \in \mathbf{R}^+$, we define $\partial_{x_k}^\alpha$ to be a pseudodifferential operator with a polyhomogeneous symbol $(2\pi i \xi_k)^\alpha$, i.e.

$$\partial_{x_k}^\alpha u = ((2\pi i \xi_k)^\alpha \hat{u}(\xi))^\vee.$$

In the sequel, we shall assume that sequences (\mathbf{u}_r) and (\mathbf{v}_r) are uniformly compactly supported. This assumption can be removed if the orders of derivatives $(\alpha_1, \dots, \alpha_d)$ are natural numbers.

Lemma. Assume that sequences (\mathbf{u}_n) and (\mathbf{v}_n) are bounded in $L^p(\mathbf{R}^d; \mathbf{R}^r)$ and $L^q(\mathbf{R}^d; \mathbf{R}^r)$, respectively, and converge toward $\mathbf{0}$ and \mathbf{v} in the sense of distributions.

Furthermore, assume that sequence (\mathbf{u}_n) satisfies:

$$\mathbf{G}_n := \sum_{k=1}^d \partial_k^{\alpha_k} (\mathbf{A}^k \mathbf{u}_n) \rightarrow \mathbf{0} \text{ in } W^{-\alpha_1, \dots, -\alpha_d; p}(\Omega; \mathbf{R}^m), \quad (2)$$

where either $\alpha_k \in \mathbf{N}$, $k = 1, \dots, d$ or $\alpha_k > d$, $k = 1, \dots, d$, and elements of matrices \mathbf{A}^k belong to $L^{\bar{s}'}(\mathbf{R}^d)$, $\bar{s} \in (1, \frac{pq}{p+q})$.

Finally, by μ denote a matrix H -distribution corresponding to subsequences of (\mathbf{u}_n) and $(\mathbf{v}_n - \mathbf{v})$. Then the following relation holds

$$\left(\sum_{k=1}^d (2\pi i \xi_k)^{\alpha_k} \mathbf{A}^k \right) \mu = \mathbf{0}.$$

Compactness by compensation result

Introduce the set

$$\Lambda_{\mathcal{D}} = \left\{ \mu \in L^{\bar{s}}(\mathbf{R}^d; (C^d(\mathbf{P}))')^r : \left(\sum_{k=1}^n (2\pi i \xi_k)^{\alpha_k} \mathbf{A}^k \right) \mu = \mathbf{0}_m \right\},$$

where the given equality is understood in the sense of $L^{\bar{s}}(\mathbf{R}^d; (C^d(\mathbf{P}))')^m$.

Let us assume that coefficients of the bilinear form q from (1) belong to space $L_{loc}^t(\mathbf{R}^d)$, where $1/t + 1/p + 1/q < 1$.

Definition. We say that set $\Lambda_{\mathcal{D}}$, bilinear form q from (1) and matrix $\mu = [\mu_1, \dots, \mu_r]$, $\mu_j \in L^{\bar{s}}(\mathbf{R}^d; (C^d(\mathbf{P}))')^r$ satisfy the strong consistency condition if $(\forall j \in \{1, \dots, r\}) \mu_j \in \Lambda_{\mathcal{D}}$, and it holds

$$\langle \phi \mathbf{Q} \otimes 1, \mu \rangle \geq 0, \quad \phi \in L^{\bar{s}}(\mathbf{R}^d; \mathbf{R}_0^+).$$

Theorem. Assume that sequences (\mathbf{u}_n) and (\mathbf{v}_n) are bounded in $L^p(\mathbf{R}^d; \mathbf{R}^r)$ and $L^q(\mathbf{R}^d; \mathbf{R}^r)$, respectively, and converge toward \mathbf{u} and \mathbf{v} in the sense of distributions.

Assume that (2) holds and that

$$q(\mathbf{x}; \mathbf{u}_n, \mathbf{v}_n) \rightharpoonup \omega \text{ in } \mathcal{D}'(\mathbf{R}^d).$$

If the set $\Lambda_{\mathcal{D}}$, the bilinear form (1), and matrix H -distribution μ , corresponding to subsequences of $(\mathbf{u}_n - \mathbf{u})$ and $(\mathbf{v}_n - \mathbf{v})$, satisfy the strong consistency condition, then

$$q(\mathbf{x}; \mathbf{u}, \mathbf{v}) \leq \omega \text{ in } \mathcal{D}'(\mathbf{R}^d).$$

H-distributions

H-distributions were introduced by N. Antonić and D. Mitrović (2011) as an extension of H-measures to the $L^p - L^q$ context.

M. Lazar and D. Mitrović (2012) extended and applied them on a velocity averaging problem.

We need multiplier operators with symbols defined on a manifold \mathbf{P} determined by an d -tuple $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbf{R}_+^d$ where $\alpha_k \in \mathbf{N}$ or $\alpha_k \geq d$:

$$\mathbf{P} = \left\{ \xi \in \mathbf{R}^d : \sum_{k=1}^d |\xi_k|^{2\alpha_k} = 1 \right\},$$

We shall use the following variant of H-distributions.

Theorem. Let (u_n) be a bounded sequence in $L^p(\mathbf{R}^d)$, $p > 1$, and let (v_n) be a bounded sequence of uniformly compactly supported functions in $L^q(\mathbf{R}^d)$, $1/q + 1/p < 1$, weakly converging to 0 in the sense of distributions. Then, after passing to a subsequence (not relabelled), for any $\bar{s} \in (1, \frac{pq}{p+q})$ there exists a continuous bilinear functional B on $L^{\bar{s}'}(\mathbf{R}^d) \otimes C^d(\mathbf{P})$ such that for every $\varphi \in L^{\bar{s}'}(\mathbf{R}^d)$ and $\psi \in C^d(\mathbf{P})$, it holds

$$B(\varphi, \psi) = \lim_{n \rightarrow \infty} \int_{\mathbf{R}^d} \varphi(\mathbf{x}) u_n(\mathbf{x}) (\mathcal{A}_{\psi} v_n)(\mathbf{x}) d\mathbf{x},$$

where \mathcal{A}_{ψ} is the Fourier multiplier operator on \mathbf{R}^d associated to $\psi \circ \pi_{\mathbf{P}}$.

The bilinear functional B can be continuously extended as a linear functional on $L^{\bar{s}'}(\mathbf{R}^d; C^d(\mathbf{P}))$.

Case $L^p - L^{p'}$, $p > 1$

In the case $1/p + 1/q = 1$, applying the same proof gives us continuous bilinear functional on $C(\mathbf{R}^d) \otimes C^d(\mathbf{P})$. Using Schwartz's kernel theorem, we can only extend it to a distribution from $\mathcal{D}'(\mathbf{R}^d \times \mathbf{P})$.

Introduce the truncation operator $T_l(v) = v$ if $|v| \leq l$ and $T_l(v) = 0$ if $|v| \geq l$, for $l \in \mathbf{N}$.

Theorem. Assume that

- sequences (\mathbf{u}_r) and (\mathbf{v}_r) are bounded in $L^p(\mathbf{R}^d; \mathbf{R}^N)$ and $L^{p'}(\mathbf{R}^d; \mathbf{R}^N)$, where $1/p + 1/p' = 1$, and converge toward \mathbf{u} and \mathbf{v} in the sense of distributions;
- for every $l \in \mathbf{N}$, the sequences $(T_l(\mathbf{v}_r))$ converge weakly in $L^{p'}(\mathbf{R}^d; \mathbf{R}^N)$ toward \mathbf{h}^l , where the truncation operator T_l is understood coordinatewise;
- there exists a vector valued function $\mathbf{V} \in L^{p'}(\mathbf{R}^d; \mathbf{R}^N)$ such that $\mathbf{v}_r \leq \mathbf{V}$ holds coordinatewise for every $r \in \mathbf{N}$;
- (2) holds with $a_{skl} \in C_0(\mathbf{R}^d)$ and that $q_{jm} \in C(\mathbf{R}^d)$.

Assume that

$$q(\mathbf{x}; \mathbf{u}_r, \mathbf{v}_r) \rightharpoonup \omega \text{ in } \mathcal{D}'(\mathbf{R}^d).$$

If for every $l \in \mathbf{N}$, the set $\Lambda_{\mathcal{D}}$, the bilinear form (1), and the (matrix of) H -distributions μ_l corresponding to the sequences $(\mathbf{u}_r - \mathbf{u})$ and $(T_l(\mathbf{v}_r) - \mathbf{h}^l)_r$ satisfy the strong consistency condition, then it holds

$$q(\mathbf{x}; \mathbf{u}, \mathbf{v}) \leq \omega \text{ in } \mathcal{D}'(\mathbf{R}^d).$$

Application

Let us consider the non-linear parabolic type equation

$$L(u) = \partial_t u - \text{div div}(g(t, \mathbf{x}, u) \mathbf{A}(t, \mathbf{x})),$$

on $(0, \infty) \times \Omega$, where Ω is an open subset of \mathbf{R}^d . For p, q and s such that $1/p + 1/q + 1/s < 1$, assume

$u \in L^p((0, \infty) \times \Omega)$, $g(t, \mathbf{x}, u(t, \mathbf{x})) \in L^q((0, \infty) \times \Omega)$,

$$\mathbf{A} \in L_{loc}^s((0, \infty) \times \Omega)^{d \times d},$$

and that the matrix \mathbf{A} is strictly positive definite, i.e.

$\mathbf{A}\xi \cdot \xi > 0$, $\xi \in \mathbf{R}^d \setminus \{0\}$, (a.e. $(t, \mathbf{x}) \in (0, \infty) \times \Omega$).

Furthermore, assume that g is a Carathéodory function and non-decreasing with respect to the third variable.

Theorem. Assume that sequences

- (u_r) and $g(\cdot, u_r)$ are such that $u_r, g(u_r) \in L^2(\mathbf{R}^+ \times \mathbf{R}^d)$ for every $r \in \mathbf{N}$;
- that they are bounded in $L^p(\mathbf{R}^+ \times \mathbf{R}^d)$, $p \in (1, 2]$, and $L^q(\mathbf{R}^+ \times \mathbf{R}^d)$, $q > 2$, respectively, where $1/p + 1/q < 1$;
- $u_r \rightharpoonup u$ and, for some, $f \in W^{-1,-2;p}(\mathbf{R}^+ \times \mathbf{R}^d)$, the sequence

$$L(u_r) = f_r \rightarrow f \text{ strongly in } W^{-1,-2;p}(\mathbf{R}^+ \times \mathbf{R}^d).$$

Under the assumptions given above, it holds

$$L(u) = f \text{ in } \mathcal{D}'(\mathbf{R}^+ \times \mathbf{R}^d).$$

References

- [1] N. Antonić, D. Mitrović: *H-distributions: An Extension of H-Measures to an $L^p - L^q$ Setting*, Abstr. Appl. Anal. (2011), Article ID 901084, 12 pages.
- [2] M. Lazar, D. Mitrović: *On an extension of a bilinear functional on $L^p(\mathbf{R}^n) \times E$ to Bôchner spaces with an application to velocity averaging*, C. R. Math. Acad. Sci. Paris 351 (2013) 261–264.
- [3] E. Yu. Panov: *Ultraparabolic H-measures and compensated compactness*, Ann. Inst. H. Poincaré Anal. Non Linéaire 28 (2011) 47–62.
- [4] L. Tartar: *H-measures, a new approach for studying homogenisation, oscillation and concentration effects in PDEs*, Proc. Roy. Soc. Edinburgh Sect. A 115 (1990) 193–230.